



UNIVERSITÉ
DE LORRAINE



paul
erlaine
université - metz
ufr sciences fondamentales
et appliquées

GRAVIER Dorian

Année 2011-2012

Rapport de stage Master Environnement et Aménagement

Spécialité GESMARE 2^{ème} année – Parcours HB

Seasonality and differences in insect emergences among
3 lakes around Constance (Germany) and their influences
on bat activity between spring and early summer.

Achieved from the 1st of March to the 31st of August 2012,
within the Limnological Institute of the Constance's
University (Mainaustraße 252, 78464 Constance –
Germany) and under the direction of:

a) Ms Ioanna **Salvarina**, Phd student within the
International Max Planck Research School (IMPRS)


And of

b) Prof. Gérard **Masson**, lecturer, teacher-researcher
within the Laboratoire des Interactions Ecotoxicologie,
Biodiversité, Ecosystèmes (LIEBE)

Thesis supported the 6th of September front of the jury
composed by Prof. Gérard Masson and Prof. Alain Izart.

Limnologisches Institut der Universität Konstanz

Universität
Konstanz



Résumé

Les échanges entre les écosystèmes aquatiques et terrestres sculptent la diversité spécifique et fonctionnelle de la nature. Les émergences d'insectes aquatiques des écosystèmes lenticques produisent d'importantes ressources saisonnières pour les consommateurs terrestres que sont les chauves-souris. Cette étude a été réalisée sur 3 lacs par l'intermédiaire de mesures hydrologiques, de piégeage d'insectes et d'enregistrements acoustiques (ultrasons). L'objectif a été d'étudier les différents liens entre la qualité de l'eau, les insectes aquatiques et terrestres et l'activité des chauves-souris, aux alentours de Constance en Allemagne. Les émergences d'insectes entre le printemps et le début d'été ont été saisonnières avec en général deux pics en termes d'intensité. L'émergence a été le plus souvent plus importante pendant les nuits. Les observations réalisées indiquent une influence combinatoire de la profondeur, de la température de l'eau, de sa qualité (comme l'eutrophisation) et des habitats alentours. L'activité des chauves-souris enregistrée provient majoritairement d'espèces opportunistes et a évolué dans le temps de la même façon que les émergences. Statistiquement, cette activité a été corrélée de façon plus importante avec le nombre d'insectes capturés sur la berge. Cependant, ce nombre a été fortement influencé par l'intensité des émergences. Les lacs étudiés diffèrent sur quasiment tous les paramètres qui les caractérisent : surface, habitats environnants, qualité de l'eau, population de poissons et de chauves-souris. Ces différences ont résulté sur l'impossibilité de prouver un réel effet de l'eutrophisation sur les variables étudiées. Cependant, cette étude aura permis de pointer les limites et les améliorations possibles de ce genre de problématique tout autant que d'émettre des hypothèses sur les échanges entre milieux aquatiques et terrestres, par le biais des émergences d'insectes aquatiques et de l'activité des chauves-souris.

Mots-clés : flux, émergence, chauves-souris, activité, lac, eutrophisation.

Abstract

Subsidies between aquatic and terrestrial systems are a driving force in the species and functional diversity in nature. Aquatic insect emergences in lentic systems are providing a seasonal and important amount of resources for terrestrial consumers as bats. This study done on 3 lakes has investigated with water measurements, insect collections and acoustic monitoring, what are the potential links between water quality, aquatic-terrestrial insects and the bat activity around Constance in Germany. Insect emergences between spring and early summer were seasonal with in general 2 peaks. The emergence was usually higher during nights. Observations indicated an influence of the combination of the depth, the water temperature, the water quality (with eutrophication) and surrounding habitats. Bat activity recorded was mainly due to opportunistic species and changed over time in the same way as the emergence. Statistically, the bat activity was more correlated with flying insects than insect emerging from lakes. However flying insects collected were highly influenced by emergences. Studied lakes are differing on almost all parameters: area, surrounding habitats, water quality, fish population and bat population. Those differences have lead on the impossibility to prove a real eutrophication effect on emergences as well as on bat activity. However this study has highlighted limits, possible improvements with such problematic and have drawn potential hypothesis on subsidies between aquatic and terrestrial systems, through aquatic insect emergences and bat activity.

Keywords: subsidies, emergence, bat activity, acoustic monitoring, lake, eutrophication.

Aquatic as well as terrestrial systems can produce material as well as organisms in various forms. This production does not always stay where it appeared and can move according to diverse forces from one system to another. Those movements are called subsidies and are known to shape, to transform, to create strong links between habitats. Subsidies are important components of the species and the functional diversity in nature. Moreover they are strongly influencing the resilience and the resistance of an entire ecosystem. They can be continuous as well as under pulse forms.

Subsidies from lentic to terrestrial systems are not extremely described. However they can have the potential to create stronger linkages than do lotic systems. The impressive example of insect emergences in some lakes is one real proof. This example is known to considerably enhance predator population living in terrestrial habitats. In Europe, insect emergences from lentic systems are mainly composed of Chironomidae (Diptera family). This family can be used as a freshwater biomonitoring.

European bats are insectivorous and can benefit from such pulse subsidies. Moreover they are really sensitive to different ecological modifications on their habitats explaining why they are considered as good bioindicators. Chironomidae emergence can be affected by different water pollution, as for example the eutrophication. This pollution is hypothetically affecting bats because depend on this food resources.

This study has the aim to increase the knowledge on aquatic insect emergences in different lakes and to study how the bat activity is affected by them. We suppose that the eutrophication as well as different other parameters can influence the seasonality of emergences, its intensity and its species diversity. Following this, the bat activity can react differently according to specificities of each species. Having better knowledge concerning all those processes can give new arguments and explanations to people who are working for lakes or bats protection.

This study is based on the collection of aquatic and terrestrial insects coupled with acoustic monitoring of bats. 3 different lakes differing in their trophic status are investigated within the protocol, followed from April until the beginning of July 2012.

The first part of this thesis is relating what is known and what is important to know on subsidies, Chironomidae and bats. The second part is commonly describing the materials and methods applied. Results are following it and are trying to give a sufficient overview on what has been observed. The last part contains a discussion describing limits on what has been done, discovered or not and is comparing results to the bibliography.

Bibliography

- Abbott I.M., Sleeman D.P. & Harrison S., 2009. Bat activity affected by sewage effluent in Irish rivers. *Biological Conservation*, 142(12), pp.2904–2914.
- Agosta S.J. & Morton D., 2003. Diet of the big brown bat, *Eptesicus fuscus*, from Pennsylvania and western Maryland. *Northeastern Naturalist*, 10(1), pp.89–104.
- Ahlen I. & Baagøe H.J., 1999. Use of ultrasound detectors for bat studies in Europe: experiences from field identification, surveys, and monitoring. *Acta Chiropterologica*, 1(2), pp.137–150.
- Aihartza J.R. et al., 2003. Evidences of piscivory by *Myotis capaccinii* (Bonaparte, 1837) in southern Iberian Peninsula. *Acta chiropterologica*, 5(2), pp.193–198.
- Akasaka T., Nakano D. & Nakamura F., 2009. Influence of prey variables, food supply, and river restoration on the foraging activity of Daubenton's bat (*Myotis daubentonii*) in the Shibetsu River, a large lowland river in Japan. *Biological Conservation*, 142(7), pp.1302–1310.
- Anderson N.H. & Cummins K.W., 1979. Influences of Diet on the Life Histories of Aquatic Insects. *Journal of the Fisheries Research Board of Canada*, 36(3), pp.335–342.
- Anderson W.B., Wait D.A. & Stapp P., 2008. Resources from another place and time: responses to pulses in a spatially subsidized system. *Ecology*, 89(3), pp.660–670.
- Anon, Times Higher Education - THE 100 Under 50 university rankings: results. Available at: <http://www.timeshighereducation.co.uk/story.asp?sectioncode=26&storycode=419908&c=1> [Accessed June 1, 2012].
- Anthony E.L.P., Stack M.H. & Kunz T.H., 1981. Night roosting and the nocturnal time budget of the little brown bat, *Myotis lucifugus*: Effects of reproductive status, prey density, and environmental conditions. *Oecologia*, 51(2), pp.151–156.
- Arthur L. & Lemaire M., 2009. Les chauves-souris de France, Belgique, Luxembourg et Suisse. *Biotope, Mèze (Collection Parthénopé)*.
- Barak Y. & Yom-Tov Y., 1989. The advantage of group hunting in Kuhl's bat *Pipistrellus kuhli* (Microchiroptera). *Journal of Zoology*, 219(4), pp.670–675.
- Barataud M., 2012. Écologie acoustique des chiroptères d'Europe: identification des espèces, étude de leurs habitats et comportements de chasse (avec DVD). *Recherche*, 67, p.02.
- Bartonicka T. & Rehak Z., 2004. Flight activity and habitat use of *Pipistrellus pygmaeus* in a floodplain forest. *Mammalia*, 68(4), pp.365–375.
- Bartonicka T. & Zukal J., 2003. Flight activity and habitat use of four bat species in a small town revealed by bat detectors. *Folia Zoologica (Brno)*, 52(2), pp.155–166.
- Basset Y., 1988. A composite interception trap for sampling arthropods in tree canopies. *Australian Journal of Entomology*, 27(3), pp.213–219.
- Baumgärtner D., Mörtl M. & Rothhaupt K.-O., 2008. Effects of water-depth and water-level fluctuations on the macroinvertebrate community structure in the littoral zone of Lake Constance. *Hydrobiologia*, 613(1), pp.97–107.
- Baxter C.V., Fausch K.D. & Carl Saunders W., 2005. Tangled webs: reciprocal flows of invertebrate prey link streams and riparian zones. *Freshwater Biology*, 50(2), pp.201–220.
- Beck A., 1995. Fecal analyses of European bat species. *Myotis*, 32(33), pp.109–119.
- Biscardi S. et al., 2007. Foraging requirements of the endangered long-fingered bat: the influence of micro-habitat structure, water quality and prey type. *Journal of Zoology*, 273(4), pp.372–381.
- Boonman M., Bretschneider F. & Van De Grind W.A., 1998. Prey detection in trawling

- insectivorous bats: duckweed affects hunting behaviour in Daubenton's bat, *Myotis daubentonii*. *Behavioral Ecology and Sociobiology*, 44(2), pp.99–107.
- Borror D.J., Triplehorn C.A. & Johnson N.F., 1989. *An introduction to the study of insects.*, Saunders College Publishing.
- Britzke E.R., 2004. Designing monitoring programs using frequency-division bat detectors: Active versus passive sampling. *Bat Echolocation Research: Tools, Techniques, and Analysis*, RM Brigham, EKV Kalko, G. Jones, S. Parsons, and HJGA Limpens, eds. Austin, TX: *Bat Conservation International*, pp.79–83.
- Broders H.G., 2003. Another quantitative measure of bat species activity and sampling intensity considerations for the design of ultrasonic monitoring studies. *Acta Chiropterologica*, 5(2), pp.235–241.
- CPEPESC Lorraine, 2009. *Connaître et protéger les Chauves-souris de Lorraine*, Ouvrage collectif coordonné par Schwaab F., Knochel A. & Jourdan D.
- Davies I.J., 1980. Relationships between dipteran emergence and phytoplankton production in the Experimental Lakes Area, northwestern Ontario. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(3), pp.523–533.
- Dechmann D.K.N. et al., 2009. Experimental evidence for group hunting via eavesdropping in echolocating bats. *Proceedings of the Royal Society B: Biological Sciences*, 276(1668), pp.2721–2728.
- Delettre Y.R. & Morvan N., 2000. Dispersal of adult aquatic Chironomidae (Diptera) in agricultural landscapes. *Freshwater Biology*, 44(3), pp.399–411.
- Dévai G. & Moldovan J., 1983. An attempt to trace eutrophication in a shallow lake (Balaton, Hungary) using chironomids. *Hydrobiologia*, 103(1), pp.169–175.
- Dickman M. & Rygiel G., 1996. Chironomid larval deformity frequencies, mortality, and diversity in heavy-metal contaminated sediments of a Canadian riverine wetland. *Environment international*, 22(6), pp.693–703.
- Dietz C. et al., 2009. *L'encyclopédie des chauves-souris d'Europe et d'Afrique du Nord: biologie, caractéristiques, protection*, Delachaux et Niestlé.
- Fenton M.B. et al., 1977. Activity patterns, habitat use, and prey selection by some African insectivorous bats. *Biotropica*, 9(2), pp.73–85.
- Fermon H. et al., 2005. Forest use and vertical stratification in fruit-feeding butterflies of Sulawesi, Indonesia: impacts for conservation. *Biodiversity and Conservation*, 14(2), pp.333–350.
- Frouz J., Ali A. & Lobinske R.J., 2002. Influence of temperature on developmental rate, wing length, and larval head capsule size of pestiferous midge *Chironomus crassicaudatus* (Diptera: Chironomidae). *Journal of economic entomology*, 95(4), pp.699–705.
- Fukui D. et al., 2006. Effect of emergent aquatic insects on bat foraging in a riparian forest. *Journal of Animal Ecology*, 75(6), pp.1252–1258.
- Füreder L., Wallinger M. & Burger R., 2005. Longitudinal and seasonal pattern of insect emergence in alpine streams. *Aquatic Ecology*, 39(1), pp.67–78.
- Gratton C., Donaldson J. & Zanden M.J., 2008. Ecosystem linkages between lakes and the surrounding terrestrial landscape in northeast Iceland. *Ecosystems*, 11(5), pp.764–774.
- Gratton C. & Zanden M.J., 2009. Flux of aquatic insect productivity to land: comparison of lentic and lotic ecosystems. *Ecology*, 90(10), pp.2689–2699.
- Greig H.S. et al., 2012. Warming, eutrophication, and predator loss amplify subsidies between aquatic and terrestrial ecosystems. *Global Change Biology*, 18(2), pp.504–514.
- Griffin D.R. & Galambos R., 1941. The sensory basis of obstacle avoidance by flying bats. *Journal of Experimental Zoology*, 86(3), pp.481–506.

- Griffin D.R., Webster F.A. & Michael C.R., 1960. The echolocation of flying insects by bats. *Animal Behaviour*, 8(3-4), pp.141–154.
- Hagen E. & Sabo J., 2011. A landscape perspective on bat foraging ecology along rivers: does channel confinement and insect availability influence the response of bats to aquatic resources in riverine landscapes? *Oecologia*, 166(3), pp.751–760.
- Hagen E.M. & Sabo J.L., 2012. Influence of river drying and insect availability on bat activity along the San Pedro River, Arizona (USA). *Journal of Arid Environments*, 84(0), pp.1–8.
- Hairston N.G., Smith F.E. & Slobodkin L.B., 1960. Community structure, population control, and competition. *American Naturalist*, pp.421–425.
- Hamilton A.L. & Saether O.A., 1971. The occurrence of characteristic deformities in the chironomid larvae of several Canadian lakes. *The Canadian Entomologist*, 103(3), pp.363–368.
- Holt R.D. & Barfield M., 2003. Impacts of temporal variation on apparent competition and coexistence in open ecosystems. *Oikos*, 101(1), pp.49–58.
- IUCN, 2008. *Red list of threatened species*, International Union for Conservation of Nature and Natural Resources Cambridge, UK.
- Jackson J.K. & Fisher S.G., 1986. Secondary Production, Emergence, and Export of Aquatic Insects of a Sonoran Desert Stream. *Ecology*, 67(3), pp.629–638.
- Jones G. & Rayner J.M.V., 1988. Flight performance, foraging tactics and echolocation in free-living Daubenton's bats *Myotis daubentoni* (Chiroptera: Vespertilionidae). *Journal of Zoology*, 215(1), pp.113–132.
- Jong J. & Ahlén I., 1991. Factors affecting the distribution pattern of bats in Uppland, central Sweden. *Ecography*, 14(2), pp.92–96.
- Kalcounis-Ruppell M.C. et al., 2007. Effects of wastewater treatment plant effluent on bat foraging ecology in an urban stream system. *Biological Conservation*, 138(1-2), pp.120–130.
- Kato C. et al., 2003. Dynamics of aquatic insect flux affects distribution of riparian web-building spiders. *Oikos*, 103(1), pp.113–120.
- Kokurewicz T., 1995. Increased population of Daubenton's bat (*Myotis daubentonii* (Kuhl, 1819))(Chiroptera: Vespertilionidae) in Poland. *Myotis*, 32(33), pp.155–161.
- Kunz T.H., 1974. Feeding Ecology of a Temperate Insectivorous Bat (*Myotis Velifer*). *Ecology*, 55(4), pp.693–711.
- Kusch J. et al., 2004. Foraging habitat preferences of bats in relation to food supply and spatial vegetation structures in a western European low mountain range forest. *FOLIA ZOOLOGICA-PRAHA*, 53(2), pp.113–128.
- Kusch J. & Idelberger S., 2005. Spatial and temporal variability of bat foraging in a western European low mountain range forest. *MAMMALIA*, 69(1), pp.21–33.
- Lacki M.J., Adam M.D. & Shoemaker L.G., 1994. Observations on Seasonal Cycle, Population Patterns and Roost Selection in Summer Colonies of *Plecotus townsendii virginianus* in Kentucky. *American Midland Naturalist*, 131(1), pp.34–42.
- Langdon P.G. et al., 2006. Assessing lake eutrophication using chironomids: understanding the nature of community response in different lake types. *Freshwater Biology*, 51(3), pp.562–577.
- Lewis T., 1970. Patterns of distribution of insects near a windbreak of tall trees. *Annals of Applied Biology*, 65(2), pp.213–220.
- Lison F. & Calvo J.F., 2011. The significance of water infrastructures for the conservation of bats in a semiarid Mediterranean landscape. *Animal Conservation*, 14(5), pp.533–541.
- Lundy M. & Montgomery I., 2009. Summer habitat associations of bats between riparian landscapes and within riparian areas. *European Journal of Wildlife Research*, 56(3), pp.385–394.
- Lynch R.J., Bunn S.E. & Catterall C.P., 2002. Adult aquatic insects: Potential contributors to

- riparian food webs in Australia's wet-dry tropics. *Austral Ecology*, 27(5), pp.515–526.
- Maier C., 1992. Activity patterns of pipistrelle bats (*Pipistrellus pipistrellus*) in Oxfordshire. *Journal of Zoology*, 228(1), pp.69–80.
- Marckmann U. & Runkel V., 2010. Automatic bat call analysis with the batcorder-system. Description of automatic bat call identification procedure and advices for the interpretation and revision of results.
- Mcaney C.M. & Fairley J.S., 1988. Activity patterns of the lesser horseshoe bat *Rhinolophus hipposideros* at summer roosts. *Journal of Zoology*, 216(2), pp.325–338.
- McCain C.M., 2007. Could temperature and water availability drive elevational species richness patterns? A global case study for bats. *Global Ecology and Biogeography*, 16(1), pp.1–13.
- McCann K., Hastings A. & Huxel G.R., 1998. Weak trophic interactions and the balance of nature. *Nature*, 395(6704), pp.794–798.
- Michelutti N. et al., 2005. Recent primary production increases in arctic lakes. *Geophys Res Lett*, 32, p.L19715.
- Mickeviciene I. & Mickevicius E., 2001. The importance of various habitat types to bats (Chiroptera: Vesperilionidae) in Lithuania during the summer period. *Acta Zoologica Lituanica*, 11(1), pp.3–14.
- Morgan N.C. & Waddell A.B., 1961. Diurnal variation in the emergence of some aquatic insects. *Transactions of the Royal Entomological Society of London*, 113(6), pp.123–137.
- Müller J. et al., 2012. Aggregative response in bats: prey abundance versus habitat. *Oecologia*. Available at: <http://www.springerlink.com/content/7n260r53u1w6388x/> [Accessed June 4, 2012].
- Nakano S. & Murakami M., 2001. Reciprocal subsidies: Dynamic interdependence between terrestrial and aquatic food webs. *Proceedings of the National Academy of Sciences*, 98(1), pp.166–170.
- Nilsson A.N., 1996a. *Aquatic insects of North Europe: a taxonomic handbook. Volume 1: Ephemeroptera, Plecoptera, Heteroptera, Neuroptera, Megaloptera, Coleoptera, Trichoptera, Lepidoptera.*, Apollo Books.
- Nilsson A.N., 1996b. *Volume 2: Aquatic insects of North Europe: a taxonomic handbook. Odonata, Diptera.*, Apollo Books.
- Norberg U.M. & Rayner J.M.V., 1987. Ecological Morphology and Flight in Bats (Mammalia; Chiroptera): Wing Adaptations, Flight Performance, Foraging Strategy and Echolocation. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 316(1179), pp.335–427.
- Nummi P. et al., 2011. Bats benefit from beavers: a facilitative link between aquatic and terrestrial food webs. *Biodiversity and Conservation*, 20(4), pp.851–859.
- O'Donnell C.F.J., 2000. Influence of season, habitat, temperature, and invertebrate availability on nocturnal activity of the New Zealand long-tailed bat (*Chalinolobus tuberculatus*). *New Zealand Journal of Zoology*, 27(3), pp.207–221.
- Ostfeld R.S. & Keesing F., 2000. Pulsed resources and community dynamics of consumers in terrestrial ecosystems. *Trends in Ecology & Evolution*, 15(6), pp.232–237.
- Paetzold A., Schubert C.J. & Tockner K., 2005. Aquatic terrestrial linkages along a braided-river: riparian arthropods feeding on aquatic insects. *Ecosystems*, 8(7), pp.748–759.
- Park K.J. & Cristinacce A., 2006. Use of sewage treatment works as foraging sites by insectivorous bats. *Animal Conservation*, 9(3), pp.259–268.
- Pinder A.M. et al., 1993. Diel periodicities of adult emergence of some chironomids (Diptera: Chironomidae) and a mayfly (Ephemeroptera: caenidae) at a western Australian wetland. *Australian Journal of Entomology*, 32(2), pp.129–135.

- Pinder L.C.V., 1986. Biology of freshwater Chironomidae. *Annual Review of Entomology*, 31(1), pp.1–23.
- Polis G.A., Anderson W.B. & Holt R.D., 1997. Toward an integration of landscape and food web ecology: the dynamics of spatially subsidized food webs. *Annual review of ecology and systematics*, pp.289–316.
- Power M.E., 1992. Top-down and bottom-up forces in food webs: do plants have primacy. *Ecology*, 73(3), pp.733–746.
- Racey P.A. & Swift S.M., 1985. Feeding Ecology of *Pipistrellus pipistrellus* (Chiroptera: Vespertilionidae) during Pregnancy and Lactation. I. Foraging Behaviour. *Journal of Animal Ecology*, 54(1), pp.205–215.
- Racey P.R. et al., 1998. Bats and insects over two Scottish rivers with contrasting nitrate status. *Animal Conservation*, 1(3), pp.195–202.
- Rosenberg D.M., 1992. Freshwater biomonitoring and Chironomidae. *Netherlands Journal of Aquatic Ecology*, 26(2-4), pp.101–122.
- Roth M., 1974. Initiation a la morphologie, la systématique et la biologie des insectes, Maurice Roth. *Initiation-Documentations techniques*. 23.
- Roué S.Y. & Barataud M., 1999. Habitats et activité de chasse des chiroptères menacés en Europe: synthèse des connaissances actuelles en vue d'une gestion conservatrice. *Le Rhinolophe*. Vol. Spec, (2), p.136.
- Runkel V., 2008. Mikrohabitatnutzung syntoper Waldfledermäuse - Usage of forest microhabitats by syntopic bat species.
- Russ J. & Montgomery W., 2002. Habitat associations of bats in Northern Ireland: implications for conservation. *Biological Conservation*, 108(1), pp.49–58.
- Russo D. & Jones G., 2003. Use of foraging habitats by bats in a Mediterranean area determined by acoustic surveys: conservation implications. *Ecography*, 26(2), pp.197–209.
- Rydell J., Entwistle A. & Racey P.A., 1996. Timing of foraging flights of three species of bats in relation to insect activity and predation risk. *Oikos*, pp.243–252.
- Rydell J., Miller L.A. & Jensen M.E., 2002. Echolocation constraints of Daubenton's Bat foraging over water. *Functional Ecology*, 13(2), pp.247–255.
- Sabo J.L. & Power M.E., 2002. River-watershed exchange: effects of riverine subsidies on riparian lizards and their terrestrial prey. *Ecology*, 83(7), pp.1860–1869.
- Saether O.A., 1979. Chironomid communities as water quality indicators. *Ecography*, 2(2), pp.65–74.
- Salvarina I. et al., 2012. Aquatic subsidies (emerging insects) to terrestrial consumers (bats). Available at: <https://www.sgmeet.com/aslo/japan2012/viewabstract2.asp?AbstractID=9716>.
- Scott S. et al., 2010. The impact of riparian habitat quality on the foraging and activity of pipistrelle bats (*Pipistrellus* spp.). *Journal of Zoology*, 280(4), pp.371–378.
- Siemers B.M. et al., 2001a. *Myotis daubentonii* is able to catch small fish. *Acta chiropterologica*, 3(1), pp.71–75.
- Siemers B.M., Stitz P. & Schnitzler H.-U., 2001b. The Acoustic Advantage of Hunting at Low Heights Above Water: Behavioural Experiments on the European “trawling” Bats *Myotis Capaccinii*, *M. Dasycneme* and *M. Daubentonii*. *Journal of Experimental Biology*, 204(22), pp.3843–3854.
- Smukalla R. & Meyer E., 1988. Insect emergence from a shallow southern West German lake, with special reference to the parasitic host-associated water mite larvae. *Hydrobiologia*, 169(2), pp.149–166.
- Stahlschmidt P. et al., 2012. Constructed wetlands support bats in agricultural landscapes. *Basic and Applied Ecology*, 13(2), pp.196–203.
- Stahlschmidt P. & Brühl C.A., 2012. Bats as bioindicators – the need of a standardized

- method for acoustic bat activity surveys. *Methods in Ecology and Evolution*. Available at: <http://onlinelibrary.wiley.com.ezproxy.ub.gu.se/doi/10.1111/j.2041-210X.2012.00188.x/full> [Accessed March 8, 2012].
- Swift S.M., Racey P.A. & Avery M.I., 1985. Feeding Ecology of *Pipistrellus pipistrellus* (Chiroptera: Vespertilionidae) During Pregnancy and Lactation. II. Diet. *Journal of Animal Ecology*, 54(1), pp.217–225.
- Taake K.H., 1991. *Strategien der Ressourcennutzung an Waldgewässern jagender Fledermäuse (Chiroptera: Vespertilionidae)*.
- Vaughan N., 1997. The diets of British bats (Chiroptera). *Mammal Review*, 27(2), pp.77–94.
- Vaughan N., Jones G. & Harris S., 1996. Effects of sewage effluent on the activity of bats (Chiroptera: Vespertilionidae) foraging along rivers. *Biological Conservation*, 78, pp.337–343.
- Vaughan N., Jones G. & Harris S., 1997. Habitat Use by Bats (Chiroptera) Assessed by Means of a Broad-Band Acoustic Method. *Journal of Applied Ecology*, 34(3), pp.716–730.
- Vermeulen A.C., 1995. Elaborating chironomid deformities as bioindicators of toxic sediment stress: the potential application of mixture toxicity concepts. In *Annales Zoologici Fennici*. pp. 265–285. Available at: <http://www.sekj.org/PDF/anzf32/anz32-265-285.pdf> [Accessed July 16, 2012].
- Voigt C.C. et al., 2011. Rain increases the energy cost of bat flight. *Biology Letters*, 7(5), pp.793–795.
- Walsh A.L. & Harris S., 1996a. Foraging habitat preferences of Vespertilionid bats in Britain. *Journal of Applied Ecology*, 33(3), pp.508–518.
- Walsh & Harris S., 1996b. Factors determining the abundance of Vespertilionid bats in Britain: geographical, land class and local habitat relationships. *Journal of Applied Ecology*, 33(3), pp.519–529.
- Walters C. et al., 2012. A continental-scale tool for acoustic identification of European bats. *Journal of Applied Ecology*.
- Wang J. et al., 2010. Seasonal variation in prey abundance influences habitat use by greater horseshoe bats (*Rhinolophus ferrumequinum*) in a temperate deciduous forest. *Canadian Journal of Zoology*, 88(3), pp.315–323.
- Ward J.V. & Stanford J.A., 1982. Thermal responses in the evolutionary ecology of aquatic insects. *Annual Review of Entomology*, 27(1), pp.97–117.
- Warren R.D. et al., 2000. The distribution of Daubenton's bats (*Myotis daubentonii*) and pipistrelle bats (*Pipistrellus pipistrellus*) (Vespertilionidae) in relation to small-scale variation in riverine habitat. *Biological Conservation*, 92(1), pp.85–91.
- Wickramasinghe L.P. et al., 2003. Bat activity and species richness on organic and conventional farms: impact of agricultural intensification. *Journal of Applied Ecology*, 40(6), pp.984–993.
- Williams K., 2012. *Bat Foraging Activity and Insect Abundance in Relation to Foraging Activity on Baylor University Campus*. Thesis. Available at: <https://beardocs.baylor.edu:8443/xmlui/handle/2104/8336> [Accessed May 10, 2012].
- Wolcott K.A. & Vulinec K., 2012. Bat Activity at Woodland/Farmland Interfaces in Central Delaware. *Northeastern Naturalist*, 19(1), pp.87–98.